







**Figure 1.** Effect on heart rate (beats per min, BPM) of intravenous administration of bupivacaine or IQB-9302 (0.1 to 10 mg/kg). Results are means of 8 anaesthetised rats for each drug.




**Figure 2a.** Effect on heart rate (beats per min, BPM) of intravenous administration of sodium chloride (control, C), bupivacaine (B), IQB-9302 (IQB), 0.1 or 0.3 mg/kg. Results are means and standard error of mean of 8 anaesthetised rats for each drug and means of 2 rats for control.





**Figure 2b.** Effect on heart rate (beats per min, BPM) of intravenous administration of sodium chloride (control, C), bupivacaine (B), IQB-9302 (IQB), 1 or 3 mg/kg. Results are means and standard error of mean of 8 anaesthetised rats for each drug and means of 2 rats for control.




**Figure 2c.** Effect on heart rate (beats per min, BPM) of intravenous administration of sodium chloride (control, C), bupivacaine (B), IQB-9302 (IQB), 10 mg/kg. Results are means and standard error of mean of 7 anaesthetised rats for bupivacaine and 6 for IQB-9302, and means of 2 rats for control.





**Figure 3.** Effect on mean arterial pressure (mmHg) of intravenous administration of bupivacaine or IQB-9302 (0.1 to 10 mg/kg). Results are means of 8 anaesthetised rats for each drug.




**Figure 4a.** Effect on mean arterial pressure (mmHg) of intravenous administration of sodium chloride (control, C), bupivacaine (B), IQB-9302 (IQB), 0.1 or 0.3 mg/kg. Results are means and standard error of mean of 8 anaesthetised rats for each drug and means of 2 rats for control.




**Figure 4b.** Effect on mean arterial pressure (mmHg) of intravenous administration of sodium chloride (control, C), bupivacaine (B), IQB-9302 (IQB), 1 or 3 mg/kg. Results are means and standard error of mean of 8 anaesthetised rats for each drug and means of 2 rats for control.





**Figure 4c.** Effect on mean arterial pressure (mmHg) of intravenous administration of sodium chloride (control, C), bupivacaine (B), IQB-9302 (IQB), 10 mg/kg. Results are means and standard error of mean of 7 anaesthetised rats for bupivacaine and 6 for IQB-9302 and means of 2 rats for control.




**Figure 5a.** Change on heart rate (beats per min, BPM) respect to predose produced by intravenous administration of sodium chloride (control, C), bupivacaine (B), IQB-9302 (IQB), 0.1 or 0.3 mg/kg. Results are means of 8 anaesthetised rats for each drug and means of 2 rats for control.





**Figure 5b.** Change on heart rate (beats per min, BPM) respect to predose produced by intravenous administration of sodium chloride (control, C), bupivacaine (B), IQB-9302 (IQB), 1 or 3 mg/kg. Results are means of 8 anaesthetised rats for each drug and means of 2 rats for control.





**Figure 5c.** Change on heart rate (beats per min, BPM) respect to predose produced by intravenous administration of sodium chloride (control, C), bupivacaine (B), IQB-9302 (IQB), 10 mg/kg. Results are means of 7 anaesthetised rats for bupivacaine, 6 for IQB-9302 and 2 for control.



**Figure 6a.** Change on mean arterial pressure (mmHg) respect to predose produced by intravenous administration of sodium chloride (control, C), bupivacaine (B), IQB-9302 (IQB), 0.1 or 0.3 mg/kg. Results are means of 8 anaesthetised rats for each drug and means of 2 rats for control.



**Figure 6b.** Change on mean arterial pressure (mmHg) respect to predose produced by intravenous administration of sodium chloride (control, C), bupivacaine (B), IQB-9302 (IQB), 1 or 3 mg/kg. Results are means of 8 anaesthetised rats for each drug and means of 2 rats for control.



**Figure 6c.** Change on mean arterial pressure (mmHg) respect to predose produced by intravenous administration of sodium chloride (control, C), bupivacaine (B), IQB-9302 (IQB), 10 mg/kg. Results are means of of 7 anaesthetised rats for bupivacaine, 6 for IQB-9302 and 2 for control.

